

B2

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 August 2003 (28.08.2003)

PCT

(10) International Publication Number
WO 03/070959 A2

(51) International Patent Classification⁷: **C12P**
(21) International Application Number: PCT/EP03/01537
(22) International Filing Date: 15 February 2003 (15.02.2003)
(25) Filing Language: English
(26) Publication Language: English

(30) Priority Data:
020 03 968.1 22 February 2002 (22.02.2002) EP

(71) Applicant (for all designated States except US): **ROCHE VITAMINS AG** [CH/CH]; 124 Grenzacherstrasse, CH-4070 Basle (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SHIMIZU, Sakayu** [JP/JP]; 6-9 Yamashita-cho, Tokiwa, Ukyo-ku, Kyoto-shi, Kyoto-fu 616-8212 (JP). **WADA, Masaru** [JP/JP]; E-204, 38-8-1 Kenjyojima, Matsuoka-cho, Yoshida-gun, Fukui-ken 910-1142 (JP).

(74) Agent: **LEDERER & KELLER**; Prinzregentenstrasse 16, 80538 München (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR PRODUCING LEVODIONE

(57) Abstract: An enone reductase characterized by a molecular mass of $61,300 \pm 5,000$ Da; NADPH and NADH as co-factor; a temperature optimum of 55-60°C at pH 7.4; a pH optimum of 4.5-8.5 and a substrate specificity on α, β -unsaturated ketones, especially derived from a yeast and a process for the preparation of levodione from ketoisophorane.

WO 03/070959 A2

- 1 -

Process for producing levodione

The present invention relates to a process for producing (6R)-2,2,6-trimethyl cyclohexane-1,4-dione (hereinafter referred to as levodione) from 2,6,6-trimethyl-2-cyclohexene-1,4-dione (hereinafter referred to as ketoisophorone) by reduced
5 nicotinamide adenine dinucleotide phosphate (herein after referred to as NADPH) dehydrogenase. Levodione is an important intermediate in the synthesis of carotenoids, e.g. zeaxanthin.

A microbiological process of producing levodione from ketoisophorone has been known, see e.g. United States Patent 4,156,100.

10 Unexpectedly, we now have found that levodione can be formed from ketoisophorone by using a NADPH dehydrogenase as a catalyst. The NADPH dehydrogenases for use as catalysts in the process of the present invention are generally known as old yellow enzyme (OYE) and defined by the enzyme class E.C. 1.6.99 according to the Enzyme Nomenclature provided by the Nomenclature
15 Committee of the International Union of Biochemistry and Molecular Biology (Academic Press, 1992).

The present invention is related to a process for producing levodione from ketoisophorone which comprises contacting ketoisophorone with NADPH dehydrogenase in the presence of NADH or NADPH in an aqueous medium, and
20 isolating the resulted levodione from the reaction mixture.

As used herein, the term "NADPH dehydrogenase" encompasses proteins capable of catalyzing the enzymatic reaction of ketoisophorone to levodione in the presence of NADH or NADPH, such as the old yellow enzyme (OYE) classified as EC 1.6.99 according to the Enzyme Nomenclature provided by the Nomenclature
25 Committee of the International Union of Biochemistry and Molecular Biology.

Particularly, the present invention is related to a process for producing

- 2 DT01 Rec'd PCT/PTC 22 MAR 2005

levodione from ketoisophorone wherein the NADPH dehydrogenase catalyzing said reaction is OYE defined by the enzyme class EC 1.6.99.

OYE for use as catalyst in the process of the present invention is obtainable or may be isolated from any appropriate microorganisms suitable for the production of said enzyme, including but not limited to the genera *Saccharomyces*, *Zygosaccharomyces*, *Candida*, *Gluconobacter*, *Beneckea*, or *Vibrio*. The above mentioned microorganisms also include synonyms or basonyms of such microorganisms having the same physico-chemical properties, as defined by the International Code of Nomenclature of Prokaryotes.

A transformed microorganism, such as *Escherichia coli*, expressing NADPH dehydrogenase can also be used as a starting microorganism.


In one embodiment of the present invention, the microorganism suitable for the production of NADPH dehydrogenase such as OYE is *Saccharomyces cerevisiae*, preferably *Saccharomyces cerevisiae* S288C (ATCC 204508) publicly available from the American Type Culture Collection, 10801 University Boulevard Manassas, VA20100-2209, U.S.A. The well known purification process can be used for the preparation of the enzyme (cf. Abramovitz, A. S., & Massey, V., J. Biol. Chem. 251,5321-5326,1976). Functional equivalents, subcultures, mutants and variants of said microorganism can also be used in the present invention.

In one embodiment of the present invention, a commercially available OYE can be used for the catalytic cleavage of ketoisophorone into levodione, for instance, NADPH-FMN Oxidoreductase® (Sigma, U.S.A.).

The OYE used preferably for the present invention is composed of two subunits, OYE2 and OYE3 having a molecular weight of 45.0 kDa and 44.9 kDa, respectively. Surprisingly, it was found that OYE2 and OYE3 not only can use NADPH, but also NADH as a co-factor for catalyzing the reaction of the present invention.

Thus, it is a further aspect of the present invention to provide a process wherein the OYE used for the catalytic reaction of ketoisophorone into levodione is encoded by the *oye2* or *oye3* gene derived from *Saccharomyces cerevisiae* S288C (ATCC 204508).

OYE of the present invention catalyzes the reduction of ketoisophorone to levodione in the presence of a co-factor according to the following formula:



For example, the standard enzyme reaction is performed as follows: the basal
5 reaction mixture (100 μ l of 1 M Tris-HCl buffer, pH 8.0, 100 μ l of 80 mM NADH,
100 μ l of 0.13 M ketoisophorone, and water to fill up to 1 ml of total volume), is
supplemented by 5 μ l of the enzyme solution, and is incubated at 20 to 40°C for 5-
300 min, preferably at 25°C for 30 min. The reaction mixture is extracted by 1 ml
of an organic solvent such as ethyl acetate, n-hexane, toluene, or n-butyl acetate to
10 recover the levodione into the organic solvent layer. The extract is analyzed by
known methods such as gas chromatography, high performance liquid
chromatography, thin layer chromatography or paper chromatography, or the like.
In case of the gas chromatography, the following conditions can be applied as one
embodiment:

15 Column: ULBON HR-20M (Shinwa, Japan) 0.25 mm ϕ x 30m
Column temperature: 160°C (constant)
Injector temperature: 250°C
Carrier gas: He (ca. 1ml/min)

The reaction can be conducted at pH values of from about 4.5 to about 8.5 in
20 the presence of NADH or NADPH in a solvent, such as Tris-HCl buffer, phosphate
buffer and the like. Preferably, the pH is between 5.0 and 8.0.

The present invention relates to a process for the production of levodione
from ketoisophorone by the help of OYE, wherein the reaction is carried out at pH
values of from 4.5 to 8.5 and at a temperature in the range of from 20 to 40°C.
25 Preferably, the reaction is carried out at pH values of from 5.0 to 8.0 and at a
temperature in the range of from 25 to 35°C.

The genes encoding the proteins OYE2 and OYE3 of the present invention
can be cloned based on the genomic sequence information of the originating
microorganism, e.g. *Saccharomyces cerevisiae*, and can be overexpressed in an
30 appropriate host organism such as *Escherichia coli*. The recombinant
microorganism, such as *Escherichia coli*, expressing NADPH dehydrogenase can be

prepared by well known recombinant technologies (cf. Molecular cloning: a Laboratory Manual, 2nd Edition / Cold Spring Harbor Laboratory Press, 1989).

Thus, the present invention concerns a process for producing levodione from ketoisophorone which comprises contacting ketoisophorone with a recombinant
5 microorganism expressing NADPH dehydrogenase or cell-free extract thereof in the presence of NADH or NADPH in an aqueous medium, and isolating the obtained levodione from the reaction mixture. Preferably, the recombinant microorganism is *Escherichia coli*.

The NADPH dehydrogenase expressed by the recombinant microorganism is
10 OYE defined by the enzyme class EC 1.6.99 as in another aspect of the present invention.

Particularly, the OYE used for the expression in the recombinant microorganism is derivable from a microorganism which is selected from the group consisting of *Saccharomyces*, *Zygosaccharomyces*, *Candida*, *Gluconobacter*,
15 *Beneckea* and *Vibrio* or functional equivalents, subcultures, mutants and variants thereof. Preferably, the OYE is derived from *Saccharomyces cerevisiae*, more preferably *Saccharomyces cerevisiae* S288C (ATCC 204508). Most preferred is OYE encoded by the *oye2* or *oye3* gene from *Saccharomyces cerevisiae* S288C (ATCC 204508).

The recombinant microorganism, such as e.g. *Escherichia coli*, expressing
20 NADPH dehydrogenase, may be cultured in a nutrient medium containing saccharides such as glucose or sucrose, alcohols such as ethanol or glycerol, fatty acids such as oleic acid, stearic acid or esters thereof, or oils such as rapeseed oil or soybean oil as carbon sources; ammonium sulfate, sodium nitrate, peptone, amino
25 acids, corn steep liquor, bran, yeast extract and the like as nitrogen sources; magnesium sulfate, sodium chloride, calcium carbonate, potassium monohydrogen phosphate, potassium dihydrogen phosphate and the like as inorganic salt sources; and malt extract, meat extract and the like as other nutrient sources. The cultivation can be carried out aerobically, normally with a cultivation period of
30 from 1 to 7 days in a medium pH of from 3.0 to 9.0 and at a temperature in the range of from 10 to 40°C. Preferably, the cultivation is carried out at the medium pH of from 5.0 and 8.0, and the cultivation temperature of from 25 to 35°C for from 2 to 4 days.

OYE is useful as a catalyst for the production of levodione from ketoisophorone. In one embodiment, the reaction for producing levodione from ketoisophorone using a recombinant microorganism can be conducted at pH values of from about 4.5 to about 8.5 in the presence of NADH or NADPH in a solvent, such as Tris-HCl buffer, phosphate buffer and the like. In a preferred embodiment, the reaction is carried out at a pH between 5.0 and 8.0.

A preferred temperature range for carrying out the reaction is from 20 to 40°C. When the pH and the temperature are set at 5.0 to 8.0 and 25 to 35°C, respectively, the reaction usually produces the best results.

In a further embodiment, the reaction for producing levodione using a recombinant microorganism is carried out at pH values of from 4.5 to 8.5 and at a temperature in the range of from 20 to 40°C. Preferably, the reaction is carried out at pH values of from 5.0 to 8.0 and at a temperature in the range of from 25 to 35°C.

The concentration of ketoisophorone in a solvent can vary depending on other reaction conditions, but, in general, is between 1 mM and 2 M, preferably between 10 mM and 100 mM.

In the reaction, OYE may also be used in an immobilized state with an appropriate carrier. Any means of immobilizing enzymes generally known in the art may be used. For instance, the enzyme may be bound directly to a membrane, granules or the like of a resin having one or more functional groups or it may be bound to the resin through bridging compounds having one or more functional groups, e.g. glutaraldehyde.

After the reaction, levodione may be recovered from the reaction mixture by extraction with an organic solvent which is non-miscible with water and which readily solubilizes levodione, such as ethyl acetate, n-hexane, toluene or n-butyl acetate. Further purification of levodione can be effected by concentrating the extract to directly crystallize levodione or by the combination of various kinds of chromatography, for example, thin layer chromatography, adsorption chromatography, ion-exchange chromatography, gel filtration chromatography or high performance liquid chromatography.

The following Examples further illustrate the present invention.

Example 1

Cloning of oye2 and oye3 genes from genomic DNA of *Saccharomyces cerevisiae*

Genomic DNA of *Saccharomyces cerevisiae* S288C (ATCC 204508) was prepared using the potassium acetate method. Using the prepared genomic DNA as template, the gene fragments of oye2 and oye3 were obtained by two-step PCR method using a thermal cycler (Perkin Elmer 2400, U.S.A.). The PCR mixture (0.02 ml) contained 5 pmol of each primer, 0.312 mM of each dNTP, and 2.5 U of Pyrobest DNA polymerase (Takara Shuzo Co. LTD / Kyoto, Japan). 100 ng of the genomic DNA of *Saccharomyces cerevisiae* S288C (ATCC 204508) was used as the template for the first PCR reaction. The mixture after the reaction was diluted 1:20 in water, and used as the template for the second PCR reaction. For the first PCR, a cycle of 10 sec. at 98°C, 30 sec. at 55°C and 90 sec. at 72°C was repeated for 25 times. For the second PCR, a cycle of 10 sec. at 98°C, 30 sec. at 51°C and 90 sec. at 72°C was repeated for 25 times.

To clone the oye2 gene, the first PCR reaction was performed with primers, OYE2-1 (5'-CGGTCCAGATATAGAATAAATCATCATATTAAG-3') (SEQ ID NO: 1), and OYE2-2 (5'-GAAATGGTGCTACAAAGTACGGTTAACAC-3') (SEQ ID NO: 2). By this reaction, DNA fragment containing the oye2 gene (1250 bp) was amplified. The second PCR was performed with primers, OYE2-3 (5'-TTAGAAGAATTCATGCCATTTGTTA-3') (SEQ ID NO: 3) and OYE2-4 (5'-AGATTTCTGCAGTTAATTTTGTCC-3') (SEQ ID NO: 4).

By this reaction, DNA fragment containing just the ORF of the oye2 gene (1200 bp) was amplified. This amplified oye2 gene was treated with *EcoRI* and *PstI*, and ligated with a vector, pKK223-3 (Amersham Bioscience / Buckinghamshire, England) that was predigested with *EcoRI* and *PstI* to construct a plasmid, pKK223-3/OYE2. *E. coli* DH5 α was transformed with the ligation mixture, and several clones were selected for sequence analysis. The sequence of the cloned oye2 gene of each candidate clone was determined by using the "Thermo Sequenase II dye terminator cycle sequencing kit" (Amersham Bioscience / Buckinghamshire, England) and an automatic sequence analyzer (ABI prism 377). Primers used for the sequence analysis were as follows :

PKK223-3(+) (5'-GACATCATAACGGTTCTGGCA-3') (SEQ ID NO: 5)

PKK223-3(-) (5'-TTATCAGACCGCTTCTGCGTT-3') (SEQ ID NO: 6)

OYE2-5 (5'-GGTATCTGGTCCGAAGAACA-3') (SEQ ID NO: 7)

OYE2-6 (5'-GACACGAGGTTCAACTAGATG-3') (SEQ ID NO: 8).

One of the clones that showed completely the same sequence as the known
oye2 sequence of *Saccharomyces cerevisiae* S288C (ATCC 204508) was selected for
further experiments. To clone the oye3 gene, the first PCR reaction was performed
with primers OYE3-1 (5'-GTACGTA CTTGATATATACAACAACTGTAG-3')
(SEQ ID NO: 9) and
OYE3-2 (5'-GCTGCCCTATATAAACAAAGATCGAGTC-3') (SEQ ID NO: 10).

By this reaction, DNA fragment containing the oye3 gene (1250 bp) was
amplified. The second PCR was performed with the following primers:

OYE3-5 (5'-TTAGAACAATTGATGCCATTTGTAA-3') (SEQ ID NO: 11)

OYE3-4 (5'-AGATTTCTGCAGTCAGTTCTTGTT-3') (SEQ ID NO: 12).

By this reaction, DNA fragment containing just the ORF of oye3 gene (1200
bp) was amplified. This amplified oye3 gene was treated with *MfeI* and *PstI*, and
ligated with a vector, pKK223-3 (Amersham Bioscience / Buckinghamshire,
England) that was predigested with *EcoRI* and *PstI* to construct a plasmid,
pKK223-3/OYE3. *E. coli* DH5 α was transformed with the ligation mixture, and
several clones were selected for sequence analysis. The sequence of the cloned oye3
gene of each candidate clone was determined by using "Thermo Sequenase II dye
terminator cycle sequencing kit" (Amersham Bioscience / Buckinghamshire,
England) and an automatic sequence analyzer (ABI prism 377). Primers used for
the sequence analysis were as follows :

PKK223-3(+) (5'-GACATCATAACGGTTCTGGCA-3') (SEQ ID NO: 13)

PKK223-3(-) (5'-TTATCAGACCGCTTCTGCGTT-3') (SEQ ID NO: 14)

OYE3-6 (5'-GACTGTGCATCTGACAGAGT-3') (SEQ ID NO: 15).

One of the clones that showed completely the same sequence as the known
oye3 sequence of *Saccharomyces cerevisiae* S288C (ATCC 204508) was selected for
further experiments.

Example 2

Levodione production using the cell-free extract of *E. coli* strain having the *oye2* or *oye3* gene of *Saccharomyces cerevisiae*

5 The plasmids, pKK223-3/OYE2 and pKK223-3/OYE3, which were constructed in the Example 1 and which comprise the complete DNA sequence of *oye2* and *oye3*, respectively, were introduced into *E. coli* JM109, and the recombinant strains, JM109[pKK223-3/OYE2] and JM109[pKK223-3/OYE3] were obtained.

10 The strain JM109[pKK223-3] was also prepared as a control. Each of these strains was inoculated into the M9 minimum medium (2 x 500 ml in 2L-Sakaguchi flask) containing 0.05 mg/ml of ampicillin and 2% (w/v) of casamino acids (Difco Laboratories, U.S.A.) and cultivated at 37°C. When the optical density at 610 nm reached 0.4, IPTG (isopropyl β -D-thiogalactopyranoside) was added to the medium to make the concentration 0.01 mM and cultivation was continued for
15 further 8-10 hours. Then the bacterial cells were collected by centrifugation. Approximately 10 g of wet cells were obtained from 1 liter of the broth. A fraction (0.7 g) of the cells was resuspended into 1.4 ml of the buffer consisting of 40 mM Tris-HCl (pH 8.0), 10 mM MgCl₂, 10 mM dithiothreitol (DTT), 200 mM KCl and 1 mM phenylmethylsulfonyl fluoride (PMSF), and the cells were disrupted with an
20 ultrasonic oscillator for 15 min. After centrifugation, the resulting supernatant was used as a cell-free extract for the levodione production as follows. Each of the cell-free extract obtained from the cells of JM109[pKK223-3/OYE2], JM109[pKK223-3/OYE3] or JM109[pKK223-3] containing 2 mg protein was used in 1 ml of the reaction mixture consisting of 25 mM Tris-HCl (pH 8.0), 66 mM NADH or 55
25 mM NADPH, and 13 mM of ketoisophorone. The reaction was carried out at 25°C for 30 minutes. The reaction mixture was extracted by 1 ml of ethylacetate to recover the levodione into ethylacetate layer. The extract was analyzed by gas chromatography [column: ULBON HR-20M (Shinwa, Japan) 0.25 mm ϕ x 30m, column temperature: 160°C (constant), injector temperature: 250°C, carrier gas:
30 He (ca. 1ml/min)]. The results are shown in Table 1.

Table 1

Clone	Co-factor	Yield of Levodione (% of ketoisophorone used)
JM109[pKK223-3/OYE2]	NADH	60
JM109[pKK223-3/OYE2]	NADPH	34
JM109[pKK223-3/OYE3]	NADH	12
JM109[pKK223-3/OYE3]	NADPH	12
JM109[pKK223-3] (control)	NADH	8.5
JM109[pKK223-3] (control)	NADPH	9.9

In another experiment using the combination of the cell-free extract of JM109[pKK223-3/OYE2] with the NADH-recycling system, the yield of levodione reached 95%. In this case, the cell-free extract of JM109[pKK223-3/OYE2] containing 30 mg protein was used in 25 ml of the reaction mixture consisting of 250 mM Tris-HCl (pH 8.0), 0.31 mM NAD⁺, 220 mM D-glucose, 12.5 units/ml glucose dehydrogenase and 65 mM of ketoisophorone. The reaction was carried out at room temperature for 90 min. The pH of the reaction mixture was controlled to be higher than 7.0 using 7% ammonium solution. As a result, 9.5 g/l of levodione (95% conversion of ketoisophorone used) was produced. Optical purity of the product was analyzed to be 94.6% (enantiomeric excess; e.e.) by gas chromatography using a chiral capillary column, BGB-176 (BGB Analytik AG, Switzerland).

Example 3

Levodione production using the cells of *E. coli* strain having the oye gene of *Saccharomyces cerevisiae*

Each of the strains, JM109[pKK223-3/OYE2] and JM109[pKK223-3], obtained in Example 2 was inoculated into the M9 minimum medium (5 ml in tube) containing 0.05 mg/ml of ampicillin and 2% (w/v) of casamino acids (Difco Laboratories, U.S.A.) and cultivated at 37°C. When the optical density at 610 nm reached 0.4, IPTG was added to the medium to make the concentration 0.01 mM and cultivation was continued for further 8-10 hours. Then the bacterial cells were collected by centrifugation, and resuspended into 2 ml of 100 mM potassium

phosphate buffer (pH 7.0). This suspension was divided into two portions (1 ml each), and the reaction was started by adding 33 mM (final concentration, hereinafter abbreviated as f.c.) of ketoisophorone and 280 mM (f.c.) of D-glucose with or without 0.37 mM (f.c.) of NAD⁺, 15 units / ml (f.c.) of glucose dehydrogenase. The reaction was carried out at 30°C overnight. The reaction mixture was extracted by ethylacetate to recover the levodione into ethylacetate layer. The extract was analyzed by gas chromatography as described in Example 1. The results are shown in Table 2.

Table 2

Clone	<u>Glucose</u> <u>dehydrogenase</u> & NAD ⁺	Yield of Levodione (% of ketoiso- phorone used)	Optical purity (% e.e.)
JM109[pKK223-3/OYE2]	-	65	59
JM109[pKK223-3/OYE2]	+	65	64
JM109[pKK223-3] (control)	-	< 1	-
JM109[pKK223-3] (control)	+	< 1	-

Claims

1. A process for producing levodione from ketoisophorone which comprises contacting ketoisophorone with NADPH dehydrogenase in the presence of NADH or NADPH in an aqueous medium, and isolating the resulted levodione from the reaction mixture.
2. The process according to claim 1, wherein the NADPH dehydrogenase is old yellow enzyme defined by the enzyme class EC 1.6.99.
3. The process according to claims 1 or 2, wherein the enzyme is obtainable from a microorganism suitable for the production of the NADPH dehydrogenase.
4. The process according to claim 3, wherein the microorganism is selected from the group of genera consisting of *Saccharomyces*, *Zygosaccharomyces*, *Candida*, *Gluconobacter*, *Beneckea*, and *Vibrio*.
5. The process according to claims 3 or 4, wherein the microorganism is *Saccharomyces cerevisiae*.
6. The process according to claims 3 or 4 wherein the microorganism is *Saccharomyces cerevisiae* S288C (ATCC 204508), a functional equivalent, subculture, mutant or variant thereof.
7. The process according to claims 1 or 2, wherein the NADPH dehydrogenase is old yellow enzyme encoded by the *oye2* or *oye3* gene derived from *Saccharomyces cerevisiae* S288C (ATCC 204508).
8. The process according to any one of claims 1 to 7, wherein the reaction is carried out at pH values of from 4.5 to 8.5 and at a temperature in the range of from 20 to 40°C.
9. The process according to any one of claims 1 to 7, wherein the reaction is carried out at pH values of from 5.0 to 8.0 and at a temperature in the range of from 25 to 35°C.
10. A process for producing levodione from ketoisophorone which

comprises contacting ketoisophorone with a transformed microorganism expressing NADPH dehydrogenase or a cell-free extract thereof in the presence of NADH or NADPH in an aqueous medium, and isolating the obtained levodione from the reaction mixture.

5 11. The process according to claim 10, wherein the recombinant microorganism is *Escherichia coli*.

12. The process according to claims 10 or 11, wherein the NADPH dehydrogenase is old yellow enzyme defined by the enzyme class EC 1.6.99.

10 13. The process according to claim 10, wherein the enzyme expressed by the transformed microorganism is derivable from a microorganism selected from the group consisting of the genera *Saccharomyces*, *Zygosaccharomyces*, *Candida*, *Gluconobacter*, *Beneckea*, and *Vibrio*.

15 14. The process according to any one of claims 10 to 13, wherein the enzyme expressed by the transformed microorganism is derived from *Saccharomyces cerevisiae*, preferably *Saccharomyces cerevisiae* S288C (ATCC 204508).

20 15. The process according to any one of claims 10 to 14, wherein the NADPH dehydrogenase expressed by the transformed microorganism is old yellow enzyme encoded by the *oye2* or *oye3* gene derived from *Saccharomyces cerevisiae* S288C (ATCC 204508).

16. The process according to any one of claims 10 to 15, wherein the reaction is carried out at pH values of from 4.5 to 8.5 and at a temperature in the range of from 20 to 40°C.

25 17. The process according to any one of claims 10 to 15, wherein the reaction is carried out at pH values of from 5.0 to 8.0 and at a temperature in the range of from 25 to 35°C.

18. The invention substantially as described hereinbefore, especially with reference to the Examples.

SEQUENCE LISTING

<110> Roche Vitamins AG
<120> Process for producing levodione
5 <130> 21147
<160> 15
<170> PatentIn version 3.1

<210> 1
10 <211> 33
<212> DNA
<213> Artificial
<400> 1

cgggtccagat atagaataaa tcatcatatt aag 33

15

<210> 2
<211> 29
<212> DNA
20 <213> Artificial

<400> 2

gaaatgggtgc tacaaagtac ggттаacac 29

25

<210> 3
<211> 25
<212> DNA
<213> Artificial
30 <400> 3

ttagaagaat tcatgccatt tgtta 25

10/528960

DT01 Rec'd PCT/PTC 22 MAR 2005

<210> 4

<211> 25

<212> DNA

5 <213> Artificial

<400> 4

agatttctgc agttaatttt tgtcc

25

10

<210> 5

<211> 21

<212> DNA

<213> Artificial

15

<400> 5

gacatcataa cggttctggc a

21

20 <210> 6

<211> 21

<212> DNA

<213> Artificial

25 <400> 6

ttatcagacc gcttctgcgt t

21

<210> 7

30 <211> 20

<212> DNA

<213> Artificial

<400> 7

ggatatctggt ccgaagaaca

20

5

<210> 8

<211> 21

<212> DNA

10 <213> Artificial

<400> 8

gacacgaggt tcaactagat g

21

15

<210> 9

<211> 30

<212> DNA

<213> Artificial

20

<400> 9

gtacgtactt gatatatata acaactgtag

30

25 <210> 10

<211> 28

<212> DNA

<213> Artificial

30 <400> 10

gctgccctat ataaacaaag atcgagtc

28

<210> 11
<211> 25
5 <212> DNA
<213> Artificial

<400> 11

ttagaacaat tgatgccatt tgtaa

25

10

<210> 12
<211> 24
<212> DNA
15 <213> Artificial

<400> 12

agattttotgc agtcagttct tggt

24

20

<210> 13
<211> 21
<212> DNA
<213> Artificial
25

<400> 13

gacatcataa cggttctggc a

21

30 <210> 14
<211> 21

<212> DNA

<213> Artificial

<400> 14

5 ttatcagacc gcttctgcgt t

21

<210> 15

<211> 20

10 <212> DNA

<213> Artificial

<400> 15

gactgtgcat ctgacagagt

20

15

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 August 2003 (28.08.2003)

PCT

(10) International Publication Number
WO 03/070959 A3

(51) International Patent Classification: C12P 7/26,
C12N 9/02, 15/53

(21) International Application Number: PCT/EP03/01537

(22) International Filing Date: 15 February 2003 (15.02.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
020 03 968.1 22 February 2002 (22.02.2002) EP

(71) Applicant (for all designated States except US): ROCHE
VITAMINS AG [CH/CH]; 124 Grenzacherstrasse,
CH-4070 Basle (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SHIMIZU, Sakayu
[JP/JP]; 6-9 Yamashita-cho, Tokiwa, Ukyo-ku, Kyoto-shi,
Kyoto-fu 616-8212 (JP). WADA, Masaru [JP/JP];
E-204, 38-8-1 Kenjyojima, Matsuoka-cho, Yoshida-gun,
Fukui-ken 910-1142 (JP).

(74) Agent: LEDERER & KELLER; Prinzregentenstrasse
16, 80538 München (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN,
YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI,
SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(88) Date of publication of the international search report:
16 October 2003

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR PRODUCING LEVODIONE

(57) Abstract: An enone reductase characterized by a molecular mass of $61,300 \pm 5,000$ Da; NADPH and NADH as co-factor; a temperature optimum of 55-60°C at pH 7.4; a pH optimum of 4.5-8.5 and a substrate specificity on α,β -unsaturated ketones, especially derived from a yeast and a process for the preparation of levodione from ketoisophorane.

WO 03/070959 A3

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/EP 03/01537

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12P7/26 C12N9/02 C12N15/53

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12P C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, FSTA, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	VAZ ALFIN D N ET AL: "Old yellow enzyme: Aromatization of cyclic enones and the mechanism of a novel dismutation reaction." BIOCHEMISTRY, vol. 34, no. 13, 1995, pages 4246-4256, XP001164161 ISSN: 0006-2960 the whole document --- -/--	1-18

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

26 August 2003

Date of mailing of the international search report

11/09/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Devijver, K

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/01537

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WANNER PETER ET AL: "Purification and characterization of two enone reductases from <i>Saccharomyces cerevisiae</i> " EUROPEAN JOURNAL OF BIOCHEMISTRY, BERLIN, DE, vol. 255, no. 1, July 1998 (1998-07), pages 271-278, XP002202649 ISSN: 0014-2956 the whole document	1-18
P,X	KATAOKA M ET AL: "OLD YELLOW ENZYME FROM <i>CANDIDA MACEDONIENSIS</i> CATALYZES THE STEREOSPECIFIC REDUCTION OF THE C = C BOND OF KETOISOPHORONE" BIOSCIENCE, BIOTECHNOLOGY AND BIOCHEMISTRY, XX, XX, vol. 66, no. 12, December 2002 (2002-12), pages 2651-2657, XP009011699 ISSN: 0916-8451 the whole document	1-18
P,X	WADA M ET AL: "PRODUCTION OF A DOUBLY CHIRAL COMPOUND, (4R,6R)-4-HYDROXY-2,2,6-TRIME THYLCYCLOHEXANONE, BY TWO-STEP ENZYMATIC ASYMMETRIC REDUCTION" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, WASHINGTON, DC, US, vol. 69, no. 2, February 2003 (2003-02), pages 933-937, XP009011700 ISSN: 0099-2240 the whole document	1-18
A	EP 1 074 630 A (HOFFMANN LA ROCHE) 7 February 2001 (2001-02-07) the whole document	1-18
A	US 4 072 715 A (BOGUTH WALTER ET AL) 7 February 1978 (1978-02-07) cited in the application the whole document	1-18
A	STOTT KELVIN ET AL: "Old Yellow Enzyme: The discovery of multiple isozymes and a family of related proteins." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 268, no. 9, 1993, pages 6097-6106, XP002252405 ISSN: 0021-9258 the whole document	1-18
	--- -/--	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 03/01537

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>NIINO YUKO S ET AL: "A new Old Yellow Enzyme of <i>Saccharomyces cerevisiae</i>." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 5, 1995, pages 1983-1991, XP002252406 ISSN: 0021-9258 the whole document</p>	1-18

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/01537

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1074630	A	07-02-2001	EP 1074630 A2	07-02-2001
			BR 0003302 A	05-06-2001
			CA 2315837 A1	02-02-2001
			CN 1285410 A	28-02-2001
			JP 2001061488 A	13-03-2001
			US 6428991 B1	06-08-2002
US 4072715	A	07-02-1978	CH 605533 A5	29-09-1978
			US 3988205 A	26-10-1976
			US 4095038 A	13-06-1978
			US 4156100 A	22-05-1979
			AT 347422 B	27-12-1978
			AT 645575 A	15-05-1978
			BE 832565 A1	20-02-1976
			DE 2537060 A1	04-03-1976
			FR 2303797 A1	08-10-1976
			FR 2303786 A1	08-10-1976
			FR 2303798 A1	08-10-1976
			GB 1508195 A	19-04-1978
			GB 1508197 A	19-04-1978
			IT 1041904 B	10-01-1980
			JP 1229051 C	19-09-1984
			JP 55033463 A	08-03-1980
			JP 59001692 B	13-01-1984
			JP 1209256 C	29-05-1984
			JP 51082789 A	20-07-1976
			JP 58007277 B	09-02-1983
			NL 7509925 A ,B,	24-02-1976
			US 4026949 A	31-05-1977
			GB 1508196 A	19-04-1978

THIS PAGE BLANK (USPTO)